

# Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.







United States  
Department of  
Agriculture

Forest Service

Pacific Northwest  
Research Station

Research Note  
PNW-RN-505  
August 1991



# *In vitro* Fruiting of *Armillaria* Species

Jimmy L. Reaves and Michael McWilliams

FEB 5 '92

USDA  
NATL. AGRIC. LIBRARY  
RECEIVED

ACQ./SERIALS BRANCH

## Abstract

Thirty-three different isolates of various *Armillaria* species were grown on sterilized orange slices under a controlled temperature and light regime. Nine isolates of *A. ostoyae* (North American Biological Species I, NABS I), three of NABS VII, one of NABS IX, one of NABS X, and two unidentified isolates formed basidiomes. Most of the basidiomes had uncharacteristic and inverted caps. Three different basidiomes contained viable spores, two from *A. ostoyae* and one from an unidentified species. Thirteen (9 NABS I, 3 NABS VII, 1 NABS X) of the identified and two of the unidentified isolates that fruited were obtained from conifers and one (NABS IX) from a hardwood. This seems to be the first report of *in vitro* production of basidiomes by NABS VII, IX, and X.

Keywords: Biological species, basidiomes, basidiocarps, sporophores.

## Introduction

Ten reproductively isolated groups (biological species<sup>1</sup>) of *Armillaria* have been identified in North America (Anderson and Ullrich 1979), five in Europe (Mohammed and others 1989), and four in Australia (Kile and Watling 1988). The pairing of haploid isolates on an agar-based medium and observing compatibility-incompatibility reactions have been the primary means of identification for these biological species (Wargo and Shaw 1985).

Identification of an *Armillaria* isolate species is reliable when known haploid isolates are paired with unknown haploid isolates, but less reliable when known haploid isolates are paired with unknown diploid isolates, such as those commonly recovered from *Armillaria*-colonized woody tissue or rhizomorphs collected *in vivo* (Siepmann 1985). Fruiting of *Armillaria* is sporadic in forests of the interior of Western North America where *Armillaria* is recognized as a serious pathogen (Wargo and Shaw 1985). If single-spore isolates are not available, species determination through haploid-diploid pairings may not clarify which *Armillaria* species are present in any particular area (Siepmann 1985).

<sup>1</sup> Biological species are groups that are reproductively isolated, and the intersterility between groups is absolute (Anderson and Ullrich 1979).

JIMMY L. REAVES is a USDA Agricultural Liaison Officer, Alabama A&M University, Normal, Alabama 35762. He was a research plant pathologist, U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, Corvallis, Oregon 97331, when research was done. MICHAEL MCWILLIAMS is a research technician, Forestry Sciences Laboratory, 3200 SW Jefferson Way, Corvallis, Oregon 97331.



*In vitro* fruitification of *Armillaria* provides a means to reduce the uncertainty of haploid-diploid pairings. Raabe (1972) successfully fruited *A. mellea* on branches of fig trees that were ground, moistened, and autoclaved, and Shaw and others (1981) were successful in getting *A. novae-zelandiae* (Stevenson) Herink. to fruit by using a defined medium. Recently, species of *Armillaria* from Africa, North America, and Europe fruited on orange slices *in vitro*<sup>2</sup> (Guillaumin and others 1989, Mohammed and others 1989, Shaw 1989).

Objectives of this study were to evaluate the feasibility of producing spore-bearing basidiocarps of various *Armillaria* species *in vitro*. Origins of the diploid isolates of *Armillaria* we used are shown in table 1.

## Materials and Methods

Four medium-sized oranges were cut into ca. 7.6-centimeter slices and placed into ca. 1-liter jars or 1-liter flasks. Distilled water was poured into the containers up to the top of the orange slices (Mohammed and others 1989). Containers were sterilized in an autoclave for 15 minutes at 15 pounds per square inch.

Containers were then seeded by placing 5-millimeter mycelial plugs (five of only one isolate in each container) of *Armillaria* from 21-day-old pure cultures of each of 32 different isolates into each container. Containers were incubated in the dark at ambient room temperature (20-23 °C) until the orange slices were thoroughly colonized by mycelia and rhizomorphs of *Armillaria* (ca. 3-4 weeks). They were placed in an environmental chamber incubator with fluorescent lights, where they were incubated under alternating 12-hour light and dark intervals at 12 °C until basidiomes developed (ca. 2-3 months).

We tried to recover single spores by selecting the best basidiome from each container, cutting the cap of the basidiome into small sections, sticking (with petroleum jelly) a section of the cap (three replications per isolate) to a petri dish top that covered a bottom filled with the enriched medium of Shaw and Roth medium (SRM) below (Shaw and Roth 1976). The petri dish tops were replaced with sterile tops after the sections of basidiomes cast spores for 1, 2, 4, or 8 hours and were incubated at 25 °C until spores germinated. A sterile needle was used to carefully remove germlings from the medium surfaces as soon as they formed. Germlings from each isolate were placed on fresh SRM and incubated in the dark at 25 °C for 21 days.

## Results and Discussion

Basidiomes formed in 16 of 33 containers (table 1). In most instances, basidiome caps were inverted, thereby revealing gills to the outside (fig. 1). Typical basidiomes developed, but in only a few containers (fig. 2). The uncharacteristic shape of the basidiome caps may have resulted from the high humidity present in the containers. In containers with typical basidiomes, there was no excess water.

<sup>2</sup> Personal communication, G. Filip, research plant pathologist, Forestry and Range Sciences Laboratory, La Grande, OR 97850.



Table 1—The production of basidiomes by *Armillaria* isolates

| <i>Armillaria</i><br>isolates | Hosts  | Location          | Year<br>isolated | Biological<br>species <sup>a</sup> | Basidiomes <sup>b</sup> | Viable<br>spores <sup>b</sup> |
|-------------------------------|--|-------------------|------------------|------------------------------------|-------------------------|-------------------------------|
| JR-D-DF-24                    | Douglas-fir<br><i>Pseudotsuga menziesii</i><br>(Mirb.) Franco)                                     | Bull Run, OR      | 1985             | NABS 1                             | +                       | —                             |
| JR-D-DF-33                    | Douglas-fir  | Glenwood, WA      | 1986             | NABS 1                             | +                       | —                             |
| Sudan #4                      | Douglas-fir  | Coast Range, OR   | 1988             | NABS 1                             | +                       | —                             |
| Sp-83-47                      | Douglas-fir  | Salmon Arm, BC    | 1983             | NABS 1                             | +                       | —                             |
| JR-D-LR-25                    | Western larch<br>( <i>Larix occidentalis</i> Nutt.)  | Sisters, OR       | 1987             | NABS 1                             | —                       | —                             |
| JR-D-PP-15                    | Ponderosa pine<br><i>Pinus ponderosa</i> Dougl.<br>ex Laws.  | Glenwood, WA      | 1972             | NABS 1                             | —                       | —                             |
| JR-D-PP-34                    | Ponderosa pine   | Glenwood, WA      | 1986             | NABS 1                             | +                       | +                             |
| JR-1953-HL                    | Western hemlock<br>( <i>Tsuga heterophylla</i><br>(Raf.) Sarg.)                                    | Glenwood, WA      | 1986             | NABS 1                             | —                       | —                             |
| JR-1953-SP                    | Sugar pine<br>( <i>P. lambertiana</i> Dougl.)  | La Pine, OR       | 1981             | NABS 1                             | —                       | —                             |
| JR-D-WF-K-4                   | White fir<br>( <i>Abies concolor</i> (Gord. &<br>Glend) Lindl. ex Hildebr.)                        | Klamath Falls, OR | 1988             | NABS 1                             | +                       | —                             |
| JR-PP                         | Ponderosa pine   | Glenwood, WA      | 1988             | NABS 1                             | +                       | +                             |
| JR-1-P                        | Ponderosa pine   | Glenwood, WA      | 1986             | NABS 1                             | +                       | —                             |
| GF-MTE-87 <sup>c</sup>        | Grand fir<br>( <i>A. grandis</i> (Dougl. ex<br>D. Don) Lindl.)                                     | La Grande, OR     | 1987             | NABS 1                             | +                       | —                             |
| SP-81-13 <sup>d</sup>         | Subalpine fir<br>( <i>A. lasiocarpa</i> (Hook.)<br>Nutt.)  | Blue River, BC    | 1981             | NABS V                             | —                       | —                             |
| SP-82-29 <sup>d</sup>         | White oak<br>( <i>Quercus garryana</i> Dougl.<br>ex Hook.)   | Victoria, BC      | 1981             | NABS VII                           | —                       | —                             |
| SP-82-72 <sup>d</sup>         | Oak ( <i>Quercus</i> ) sp.   | Victoria, BC      | 1981             | NABS VII                           | —                       | —                             |
| JR-D-WF-1                     | White fir  | Weed, CA          | 1985             | NABS VII                           | —                       | —                             |
| JR-D-WF-9                     | White fir  | Mount Shasta, CA  | 1985             | NABS VII                           | —                       | —                             |
| BR-8                          | Douglas-fir  | Bull Run, OR      | 1987             | NABS VII                           | +                       | —                             |
| BR-9                          | Douglas-fir  | Bull Run, OR      | 1987             | NABS VII                           | +                       | —                             |
| JR-D-SRF-K3                   | California red fir<br>( <i>A. magnifica</i> A. Murr.)  | Klamath Falls, OR | 1988             | NABS VII                           | +                       | —                             |
| JR-D-OK-5 <sup>e</sup>        | Shingle oak<br>( <i>Q. imbricaria</i> Michx.)  | Pennsylvania      |                  | NABS IX                            | +                       | —                             |
| SP-84-15 <sup>d</sup>         | Birch<br>( <i>Betula</i> sp. Laws.)  | Fernie, BC        | 1984             | NABS X                             | —                       | —                             |
| GF-MC-87 <sup>c</sup>         | Grand fir  | Cove, OR          | 1987             | NABS X                             | +                       | —                             |
| Tree 4674                     | Douglas-fir  | Bull Run, OR      | 1985             | NA                                 | +                       | —                             |
| JR-D-CD-20                    | Western red-cedar<br>( <i>Thuja plicata</i><br>Donn. ex D. Don.)                                   | Tillamook, OR     | 1987             | NA                                 | —                       | —                             |
| JR-D-LP                       | Sierra lodgepole pine<br>( <i>P. contorta</i> var.<br><i>murrayana</i> (Grev. &<br>Balf.) Engelm.) | Sisters, OR       | 1986             | NA                                 | —                       | —                             |



Table 1—continued

| <i>Armillaria</i><br>isolates | Hosts   | Location          | Year<br>isolated | Biological<br>species <sup>a</sup> | Basidiomes <sup>b</sup> | Viable<br>spores <sup>b</sup> |
|-------------------------------|---|-------------------|------------------|------------------------------------|-------------------------|-------------------------------|
| JR-D-PP-K2                    | Ponderosa pine  | Klamath Falls, OR | 1988             | NA                                 | —                       | —                             |
| JR-D-OK0-27                   | Oregon white oak  | Lebanon, OR       | 1985             | NA                                 | —                       | —                             |
| GF-KF-77 <sup>c</sup>         | Incense cedar<br>( <i>Librocedrus decurrens</i><br>Torr.) | Klamath Falls, OR | 1977             | NA                                 | —                       | —                             |
| GF-LS-88 <sup>c</sup>         | Grand fir   | Elgin, OR         | 1988             | NA                                 | —                       | —                             |
| GF-S-79 <sup>c</sup>          | Ponderosa pine  | Sisters, OR       | 1979             | NA                                 | +                       | +                             |

NA = not available.

<sup>a</sup> Biological species were determined as described by Morrison and others (1985). NABS = North American Biological Species.

<sup>b</sup> Plus (+) = present; minus (—) = absent.

<sup>c</sup> *Armillaria* isolates obtained from Gregory Filip, USDA Forest Service, Pacific Northwest Research Station, La Grande, Oregon.

<sup>d</sup> *Armillaria* isolates obtained from Duncan Morrison, Canadian Forestry Service, Pacific Forestry Centre, Victoria, British Columbia.

<sup>e</sup> *Armillaria* isolates obtained from Philip Wargo, USDA Forest Service, Northeast Forest Experiment Station, Hamden, Connecticut.




Figure 1 (left)—Basidiomes (arrowheads) of NABS VII (BR-8) forming on the surface of orange slices. The caps are turned inside out thereby exposing the gills and the curved stalks.

Figure 2 (right)—Basidiome (arrowhead) of NABS I (JR-D-WF-K-4) forming on the surface of orange slices. The cap and stalk are characteristic.

Viable spores were recovered from only three fruiting bodies, two from NABS I isolates of and one from an unidentified species. Of those basidiomes yielding single spores, one had characteristic morphology and two had uncharacteristic morphology. Perhaps some basidiomes were infertile, or basidiomes with the exposed gills cast their spores as soon as they opened. Consequently, when we tried to obtain single spores, the gills were depleted.



Nine of the isolates that produced basidiomes were NABS I (*A. ostoyae* [Romang.] Herink.), three were NABS VII (*A. bulbosa* (Barla) Kile et Watling), one was NABS IX, one was NABS X, and two were from unidentified species (table 1). Of the 16 *Armillaria* isolates that produced basidiomes, 15 were isolated from conifers and one from a hardwood (table 1). To our knowledge, this is the first report of successful *in vitro* fruiting of known isolates of NABS VII, IX, and X of *Armillaria*. We have no definitive reason as to why NABS VII, IX, and X fruited. We did notice, however, that if the containers were not exposed to adequate light (they were shadowed by other containers), fruiting was absent or delayed. Also, in the containers where the orange slices were not completely colonized by *Armillaria* and excess water remained in the container, basidiomes did not form, perhaps because of near or actual anaerobic conditions.

If this method can be refined so that fertile basidiomes can be produced reliably by different *Armillaria* species and single spores can be consistently recovered from them, then we may be able to more accurately identify field collected isolates of *Armillaria* (Wargo and Shaw 1985).

## Acknowledgments

The authors thank Philip M. Wargo, USDA Forest Service, Northeast Forest Experiment Station, Hamden, Connecticut; Duncan J. Morrison, Canadian Forestry Service, Pacific Forestry Centre, Victoria, British Columbia; and Gregory Filip, USDA Forest Service, La Grande, Oregon, for providing some of the isolates used in this study; and Harlan Fay for his technical assistance.


## Literature Cited

- Anderson, James B.; Ullrich, Robert C. 1979. Biological species of *Armillaria mellea* in North America. *Mycologia*. 71: 402-414.
- Guillaumin, Jean-Jacques; Mohammed, Caroline; Berthelay, Susanne. 1989. *Armillaria* species in the northern temperate hemisphere. In: Morrison, Duncan J., ed. The proceedings of the 7th international conference on root and butt rots of forest trees, IUFRO; 1988 August 9-16; Vernon and Victoria, BC. [Location of publisher unknown]: [Publisher unknown]: 27-44.
- Kile, Glen A.; Watling, Roy. 1988. Identification and occurrence of Australian *Armillaria* species, including *A. pallidula* sp. nov. and comparative studies between them and non-Australian tropica and Indian *Armillaria*. *Transactions of the British Mycological Society*. 91: 305-315.
- Mohammed, Caroline; Guillaumin, Jean-Jacques; Berthelay, Susanne. 1989. Preliminary investigations about taxonomy and genetics of African *Armillaria* species. In: Morrison, Duncan J., ed. The proceedings of the 7th international conference on root and butt rots of forest trees, IUFRO; 1988 August 9-16; Vernon and Victoria, BC. [Location of publisher unknown]: [Publisher unknown]: 447-457.
- Morrison, Duncan J.; Chu, Dorothy; Johnson, Al L.S. 1985. Species of *Armillaria* in British Columbia. *Canadian Journal of Plant Pathology*. 7: 242-246.
- Raabe, Robert D. 1972. Variation in pathogenicity and virulence in single-spore isolates of *Armillaria mellea*. *Mycologia*. 64: 1154-1159.



- Shaw, Charles G., III. 1989.** *Armillaria ostoyae* associated with mortality of new hosts in Chihuahua, Mexico. *Plant Disease*. 73: 75.
- Shaw, Charles G., III; Roth, Lewis F. 1976.** Persistence and distribution of an *Armillaria* clone in a ponderosa pine forest. *Phytopathology*. 66: 1210-1213.
- Shaw, Charles G., III; Mackenzie, Martin; Toes, Elizabeth A.H.; Hood, Ian A. 1981.** Cultural characteristics and pathogenicity to *Pinus radiata* of *Armillaria novae-zelandiae* and *A. limonea*. *New Zealand Journal of Forestry Science*. 11: 6-70.
- Siepmann, Von R. 1985.** Über das Vorkommen von *Armillaria*-Arten und-klonen in benachbarter Koniferenbeständen, Mischbeständen in Laubwald. *European Journal Forest Pathology*. 15: 71-80.
- Wargo, Phillip M.; Shaw, Charles G., III. 1985.** *Armillaria* root rot: the puzzle is being solved. *Plant Disease*. 69: 826-832.





The **Forest Service** of the U.S. Department of Agriculture is dedicated to the principle of multiple use management of the Nation's forest resources for sustained yields of wood, water, forage, wildlife, and recreation. Through forestry research, cooperation with the States and private forest owners, and management of the National Forests and National Grasslands, it strives—as directed by Congress—to provide increasingly greater service to a growing Nation.

The U.S. Department of Agriculture is an Equal Opportunity Employer. Applicants for all Department programs will be given equal consideration without regard to age, race, color, sex, religion, or national origin.

Pacific Northwest Research Station  
319 S.W. Pine St.  
P.O. Box 3890  
Portland, Oregon 97208-3890





U.S. Department of Agriculture  
Forest Service

Forest Management Plan  
Forest Management Plan

Forest Management Plan  
Forest Management Plan

Forest Management Plan  
Forest Management Plan

Forest Management Plan  
Forest Management Plan

Forest Management Plan  
Forest Management Plan

Forest Management Plan  
Forest Management Plan

Forest Management Plan  
Forest Management Plan

Forest Management Plan  
Forest Management Plan

Forest Management Plan  
Forest Management Plan

Forest Management Plan  
Forest Management Plan

Forest Management Plan  
Forest Management Plan

Forest Management Plan  
Forest Management Plan

Forest Management Plan  
Forest Management Plan

Forest Management Plan  
Forest Management Plan

Forest Management Plan  
Forest Management Plan

Forest Management Plan  
Forest Management Plan

Forest Management Plan  
Forest Management Plan

Forest Management Plan  
Forest Management Plan

Forest Management Plan  
Forest Management Plan

BULK RATE  
POSTAGE  
FEES PAID  
USDA-FS  
PERMIT No.

U.S. Department of Agriculture  
Pacific Northwest Research Station  
319 S.W. Pine Street  
P.O. Box 3890  
Portland, Oregon 97208

Official Business  
Penalty for Private Use, \$300

do NOT detach label